Prepared for submission to JCAP

A Near Horizon Stress Tensor for Blackfold

arXiv:1601.07467v1 [hep-th] 27 Jan 2016

J. Sadeghi,[a] **Z. amoozad,**[a,1]

[a]University of Mazandaran,
Babolsar, Iran

E-mail: pouriya@ipm.ir, z.amoozad@stu.umz.ac.ir

Abstract. In literature, it is nicely proved that Einstein's field equations could be written near a horizon as a thermodynamical equation for the 4 dimensional spherically symmetric static and stationary spacetimes and also axisymmetric cases. In addition, for few types of higher dimensional black branes this thermodynamical equation has been derived. We determine the stress energy tensor for another interesting higher dimensional black branes which are known as blackfolds. Computations are done for two cases of charged and neutral blackfols. The method used is the near horizon expansion of the metrics of neutral and charged blackfolds. Using some rules and conditions, thermodynamic quantities of blackfolds have been calculated. Results surprisingly show that temperature, entropy, pressure and energy density are proportional to n and Ω_{n+1}. Although temperatures (and entropies) differ from each other, the obtained stress energy tensor is the same for neutral and charged blackfolds. It proves that local distribution of energy is the same for the effective theory of charged and neutral blackfolds. In addition, charged and neutral blackfolds can be regarded as a perfect fluid and requiring the general covariance for the intrinsic fluctuations, the hydrodynamic Euler equation will be reproduced.

[1]Corresponding author.

Contents

1 Introduction **1**

2 General properties of blackfolds **2**
2.1 Neutral blackfolds 3
2.2 Charged blackfolds 3

3 Near horizon expansion of neutral blackfold **4**

4 Near horizon expansion of charged blackfolds **6**

5 Conclusion and outlook **7**

1 Introduction

Investigation of stress energy tensor in hydrodynamics and fluid dynamic theories is very familiar and interesting for physicists. This quantity is a diagonal tensor which it's dimension is the dimension of the corresponding spacetime. The role and appearance of this quantity in high energy physics is one of the most intriguing concepts which attracts much attentions. In curved spacetime, and consequently in higher dimension paradigm stress energy tensor is a local quantity in contrast to the flat spacetime quantum field theory in which the physical quantities are global. The point- dependency of physical observables has been best expressed in the Einstein's field equations in $T_{\mu\nu}(x)$, $R_{\mu\nu}(x)$.

Using similar known techniques and definitions in 4 dimensional black holes, thermodynamical quantities of higher dimensional black holes have been determined by some nice methods [1–5]. For the spherically symmetric static and stationary spacetimes, and also for axisymmetric ones, it has been proved that Einstein's field equations near horizon can be written as a thermodynamical equation from which the temperature, entropy, pressure, and energy density of the black hole could be determined.

Another method which is useful for finding quasilocal stress tensor of field theory on the boundary of gravity is the well known Brown-York method [6, 7]. In this method the tensor of dual theory could be extracted by foliation of spacetime into hypersurfaces and then by calculating the extrinsic curvature, the quasilocal stress tensor takes the form:

$$T_{ab} = \frac{1}{8\pi G}(k_{ab} - \gamma_{ab}k), \tag{1.1}$$

where k_{ab} is the extrinsic curvature tensor, k is it's trace and γ_{ab} is the induced metric on that hypersurface. But in AdS spacetime (which is especial cases and important for physicists) the boundary metric takes an infinite Weyl factor as $r \longrightarrow \infty$. So this tensor can not be the correct stress tensor on the boundary. To overcome this problem and cancel divergencies, some counterterms must be added to the action, so the quasilocal stress tensor reads [8]

$$T_{ab} = \frac{1}{8\pi G}(k_{ab} - \gamma_{ab}k + \frac{2}{\sqrt{-\gamma}}\frac{\delta S_{ct}}{\delta \gamma^{ab}}), \tag{1.2}$$

which S_{ct} depends on the dimension of AdS spacetime and must be added to extract the correct boundary stress tensor.

Another notable appearance of stress energy tensor is in field theories which lives in holographic principle. As may be known, because of the existence of some kind of divergencies in field theories, it is hard to determine correct stress energy tensor for them. But there is a surprising duality which relates the calculation in gravity side to that in field theory side. This is gauge theory/gravity correspondence which for the case of AdS gravity has exactly proved by Maldacena [9–12]. It is notable that some new works have been emerged on asymptotic flat spacetimes [13–15].

The discussion shows the importance and applicational meaning of the stress tensor. As mentioned, for all of the spherically and axisymmetric 4 dimensional spacetimes, corresponding thermodynamic properties could be derived by expanding the Einstein equations near the horizon. This method shows elegantly that black hole's horizon is a thermodynamical system.

Reviewing the methods discussed above, we extend the near horizon method to the blackfolds as new higher dimensional objects.

As we know, the field theory in d+1 dimensions correspond to a statistical mechanics if the (d+1)'th dimension be periodic. After some computations, the temperature of the blackfold will be obtained. The entropy, pressure and energy density of blackfold will be extracted. Then the stress tensor of the charged and neutral blackfold will be determined. As the metric of the neutral blackfold is Ricci-flat and as recently an interesting connection has been established between AdS and Ricci-flat spacetimes [16, 17], the AdS form of the blackfold will be reproduced. The results show that the stress tensor is exactly the same as that for the AdS and Ricci-flat form of the metric of the nutral blackfold. But in the charged case, it's metric can not be transform to the AdS form because it is not Ricci-flat. Such a comparison does not exist for the charged blackfold and just the stress energy tensor for the given metric is extracted by the method of near horizon expansion. The paper is organized as follows. In the next section some properties and metrics of neutral and charged blackfolds have been reviewed. In section 3 , 4 the near horizon expansion of neutral and charged blackfold and it's stress tensor is calculated respectively. In the final section we conclude and note some points.

2 General properties of blackfolds

In high energy physics and string theory, introducing new dimensions (more than 4) often solves some fundamental problems. Extensions of Schwarzschild black hole to higher dimensions by Kaluza-Klein theory and introducing one or more spatial extra dimensions is a simple example with interesting properties [7]. If the worldvolume of a black p-brane bent into the shape of a compact hypersurface, for instance that of a torus or a sphere, many new geometries and topologies of black hole horizon would obtain. For cases which worldvolume of black brane is not exactly flat, or not in stationary equilibrium, in the way that deviation from the flat stationary black brane exists on scales much longer than the brane thickness, an effective theory could be defined. Black branes whose worldvolume is bent into the shape of a submanifold of a background spacetime have been named blackfolds [18–21]. Blackfolds have two length scales which are given by,

$$l_M \sim (GM)^{1/D-3} \quad , \quad l_J \sim \frac{J}{M}. \tag{2.1}$$

In case which $l_J \gg l_M$, this separation suggests an effective description of long wavelength dynamics. Also it may causes to the Gregory-Laflamme instability of blackfolds.

As the theory of classical brane dynamics explains a long wavelength effective theory, black branes can also take the form of a dynamic fluid that lives on a dynamical worldvolume. As we know there is a duality between Ricci-flat and AdS spacetime; using the AdS/Ricci-flat correspondence and neutral blackfold as a Ricci-flat case, the AAdS form of a neutral blackfold in the Fefferman-Graham coordinate has been extracted and the dual renormalized stress tensor which is a property of the boundary have been obtained [22].

2.1 Neutral blackfolds

As mentioned before, blackfolds can have $\frac{l_M}{l_J} \to 0$, so that a flat black brane produces and effective theory describes the collective dynamics of a black p-brane where it's geometry in $D = n + p + 3$ spacetime dimension is,

$$ds^2{}_{p-brane} = -(1 - \frac{{r_0}^n}{r^n})dt^2 + \sum_{i=1}^{p} {dz_i}^2 + (1 - \frac{{r_0}^n}{r^n})^{-1} dr^2 + r^2 d\Omega^2{}_{n+1}, \tag{2.2}$$

where r_0 is the thickness of the horizon and $\sigma^a = (t, z^i)$ is related to brane worldvolume. Here $(p + 1)$ coordinates are on the worldvolume of the blackfold and $(D - p - 1)$ coordinates are transverse directions to the worldvolume.

For isotropic worldvolume theory, in case of lowest derivative order, the stress tensor will be the same as the isotropic perfect fluid, which is given by,

$$T^{ab} = (\varepsilon + p)u^a u^b + p\gamma^{ab}, \tag{2.3}$$

where ε is the energy density and p is the pressure.

2.2 Charged blackfolds

In supergravity and low energy limit of string theory, black branes have charges. One of the best choices is black p-branes that carry charges of Ramond-Ramond field strength type $F_{(p+2)}$. The charged dilatonic black p-brane solution of action is,

$$I = \frac{1}{16\pi G} \int dx^D \sqrt{-g}(R - \frac{1}{2}(\partial\phi)^2 - \frac{1}{2(p+2)!} e^{a\phi} F^2_{(p+2)}), \tag{2.4}$$

where

$$a^2 = \frac{4}{N} - \frac{2(p+1)n}{D-2} \qquad , \qquad n = D - p - 3. \tag{2.5}$$

The flat black p-brane solution reads,

$$ds^2 = H^{\frac{-Nn}{D-2}}(-f dt^2 + \sum_{i=1}^{p} {dz_i}^2) + H^{\frac{N(p+1)}{D-2}}(f^{-1}dr^2 + r^2 d\Omega^2{}_{n+1}), \tag{2.6}$$

$$\begin{aligned} &e^{2\phi} = H^{aN} \quad , \quad A_{p+1} = \sqrt{N}\coth\alpha(H^{-1} - 1)dt \wedge dz_1 \wedge dz_2 ... \wedge dz_p, \\ &H = 1 + \frac{{r_0}^n \sinh^2\alpha}{r^n} \qquad , \qquad f = 1 - \frac{{r_0}^n}{r^n}. \end{aligned} \tag{2.7}$$

where $\sinh\alpha$ is related to the boost of the blackfold which is a Lorentz transformation of some of directions of p-brane. As $a^2 \geq 0$, the parameter N is not arbitrary and there is,

$$N \leq 2(\frac{1}{n} + \frac{1}{p+1}). \tag{2.8}$$

In string/M theory, N is an integer up to 3 (when $p \geq 1$)that corresponds to the number of different types of branes in an intersection. In this case the effective stress tensor can be written as,

$$T^{ab} = \tau s(u^a u^b - \frac{a}{b}\gamma^{ab}) - \Phi_p Q_p \gamma_{ab}, \tag{2.9}$$

where τ is temperature and s is entropy and Φ_p is the potential that measured from the difference between the values of A_{p+1} at the horizon at $r \to \infty$ in (2.7).

3 Near horizon expansion of neutral blackfold

The case of neutral blackfold has been reviewed in the section 2 and the appropriate metric is

$$ds^2{}_{p-brane} = -(1 - \frac{{r_0}^n}{r^n})dt^2 + \sum_{i=1}^{p} {dz_i}^2 + (1 - \frac{{r_0}^n}{r^n})^{-1} dr^2 + r^2 d\Omega^2{}_{n+1}. \tag{3.1}$$

In this method we must expand the metric around the horizon, so by considering $r \to r_0 + u$ and $u \ll r_0$, we have

$$\lim_{r \to r_0 + u} (1 - \frac{{r_0}^n}{r^n}) = \frac{nu}{r_0}, \tag{3.2}$$

so the metric takes the form of

$$ds^2{}_{p-brane} = -(\frac{nu}{r_0})dt^2 + \sum_{i=1}^{p} {dz_i}^2 + \frac{r_0}{nu} du^2 + {r_0}^2 d\Omega^2{}_{n+1}, \tag{3.3}$$

because the Euclidian quantum field theory in $d+1$ dimensions is equivalent to statistical mechanics in d dimensions (if the $(d+1)$'th dimension be periodic [23]), so

$$dr^2 \to du^2, \qquad t = \tau \to -it, \tag{3.4}$$

$$ds^2{}_{p-brane} = -(\frac{nu}{r_0})d\tau^2 + \sum_{i=1}^{p} {dz_i}^2 + \frac{r_0}{nu} du^2 + {r_0}^2 d\Omega^2{}_{n+1}, \tag{3.5}$$

by changing the variables,

$$\frac{du^2}{u} = d\rho^2, \qquad \to \qquad \rho = 2\sqrt{u}, \tag{3.6}$$

then

$$ds^2{}_{p-brane} = \frac{r_0}{n}\{\frac{n^2\rho^2}{4{r_0}^2} d\tau^2 + d\rho^2\} + \sum_{i=1}^{p} {dz_i}^2 + {r_0}^2 d\Omega^2{}_{n+1}, \tag{3.7}$$

by changing the variable $\theta = \frac{n\tau}{2r_0}$ then we can rewrite,

$$ds^2{}_{p-brane} = \frac{r_0}{n}\{\rho^2 d\theta^2 + d\rho^2\} + \sum_{i=1}^{p} {dz_i}^2 + {r_0}^2 d\Omega^2{}_{n+1}, \tag{3.8}$$

as we can take the following:

$$\tau \to \tau + \beta \qquad \Rightarrow \qquad \frac{n}{2r_0}\tau \to \frac{n}{2r_0}\tau + \frac{n}{2r_0}\beta, \tag{3.9}$$

and

$$\theta \to \theta + 2\pi, \tag{3.10}$$

so one can easily find

$$\frac{n}{2r_0}\beta = 2\pi \ , \quad T = \frac{1}{\beta} \quad \Rightarrow \quad T = \frac{n}{4r_0\pi}. \tag{3.11}$$

Now, we can determine the entropy of blackfold by knowing that at any given point on the worldvolume the Bekenestein-Hawking identification between horizon area and entropy obtained by compactifying the p directions along the brane. Thus the entropy density s takes the form,

$$s = \frac{A}{4G} = \frac{r_0^{n+1}\Omega_{n+1}}{4G}. \tag{3.12}$$

After finding s we can determine P as,

$$dP = sdT \qquad P = -\frac{\Omega_{n+1}r_0^n}{16\pi G}, \tag{3.13}$$

and by the relation $d\varepsilon = Tds$ the energy density of that is

$$\varepsilon = \frac{\Omega_{n+1}}{16\pi G}r_0^n, \tag{3.14}$$

so obviously we prove that the thermodynamic relations are correct and

$$\varepsilon + P = Ts. \tag{3.15}$$

The stress energy tensor of neutral blackfold takes the following form

$$T_{ab} = (\epsilon + P)u_a u_b + P\eta_{ab}, \tag{3.16}$$

then

$$T_{ab} = \frac{\Omega_{n+1}r_0^n}{16\pi G}(nu_a u_b - \eta_{ab}). \tag{3.17}$$

We have exactly found the stress tensor which we expected. This tensor is conserved and traceless. It can be obtained from an ADM-type prescription [24],or equivalently, from the Brown-York quasilocal stress-energy tensor [6]. equation (3.17), based on general covariance for the intrinsic fluctuations, obeys the relation

$$D_a T^{ab} = 0, \tag{3.18}$$

where D_a is the covariant derivative for the worldvolume metric [19]. Since the extracted stress tensor could be regarded as the stress energy tensor of a perfect fluid, the above constraint is hydrodynamic Euler equation.

Now by using the method of AdS/Ricci-flat correspondence the AdS form of the neutral blackfold can be reproduced [22],

$$ds^2{}_\Lambda = -\frac{1}{r^2}(1 - \frac{r^d}{{r_0}^d})dt^2 + \frac{1}{r^2}(dz_i{}^2 + d\vec{y}^2) + \frac{1}{r^2(1 - \frac{r^d}{{r_0}^d})}dr^2. \tag{3.19}$$

By expanding this metric near horizon and using the same method we conclude,

$$T = \frac{-d}{4\pi r_0}, \tag{3.20}$$

as in the transformation from AdS to Ricci-flat we must change $n \to -d$.

4 Near horizon expansion of charged blackfolds

For the case of charged dilatonic blackfold (2.6) we can repeat the method. The metric of interest is:

$$ds^2{}_{p-brane} = Adt^2 + Bdz_i{}^2 + Cdr^2 + F_\alpha d\psi^2{}_\alpha, \tag{4.1}$$

and

$$\begin{aligned} A &= -(1+\frac{{r_0}^n \sinh^2\alpha}{r^n})^{\frac{-Nn}{D-2}}(1-\frac{{r_0}^n}{r^n}),\\ B &= (1+\frac{{r_0}^n \sinh^2\alpha}{r^n})^{\frac{-Nn}{D-2}},\\ C &= (1+\frac{{r_0}^n \sinh^2\alpha}{r^n})^{\frac{N(p+1)}{D-2}}(1-\frac{{r_0}^n}{r^n})^{-1},\\ F_\alpha &= F_\alpha(r,\psi_\alpha). \end{aligned} \tag{4.2}$$

Now we try to expand the metric around horizon; $r \to r_0 + u$ and $u \ll r_0$;

$$ds^2 = -(1+\sinh^2\alpha)^{-\frac{n}{8}}\frac{nu}{r_0}dt^2+(1+\sinh^2\alpha)^{-\frac{n}{8}}dz_i{}^2+(1+\sinh^2\alpha)^{-\frac{8-n}{8}}\frac{r_0}{nu}dr^2+(1+\sinh^2\alpha)^{-\frac{8-n}{8}}d\Omega^2{}_{n+1}. \tag{4.3}$$

Again by changing the variable that we proceed in the previous section we obtain;

$$ds^2 = -(1+\sinh^2\alpha)^{-\frac{8-n}{8}}\frac{r_0}{n}\{(1+\sinh^2\alpha)^{-1}\frac{n^2 4{r_0^2}^2}{\rho}\,d\tau^2+d\rho^2\}+(1+\sinh^2\alpha)^{-\frac{n}{8}}dz_i{}^2+(1+\sinh^2\alpha)^{-\frac{8-n}{8}}d\Omega^2{}_{n+1}, \tag{4.4}$$

in which $\theta = (1+\sinh^2\alpha)^{-\frac{1}{2}}\frac{n}{2r_0}\tau$ parameters must obey:

$$\tau \to \tau+\beta \quad \Rightarrow \quad (1+\sinh^2\alpha)^{-\frac{1}{2}}\frac{n}{2r_0}\tau \to (1+\sinh^2\alpha)^{-\frac{1}{2}}\frac{n}{2r_0}\tau + (1+\sinh^2\alpha)^{-\frac{1}{2}}\frac{n}{2r_0}\beta, \tag{4.5}$$

$$\theta \to \theta+2\pi \quad \Rightarrow \quad (1+\sinh^2\alpha)^{-\frac{1}{2}}\frac{n}{2r_0}\tau \to (1+\sinh^2\alpha)^{-\frac{1}{2}}\frac{n}{2r_0}\tau + 2\pi, \tag{4.6}$$

so the temperature of blackfold takes the form,

$$T = \frac{1}{\beta} = \frac{n}{4\pi r_0}(1+\sinh^2\alpha)^{-\frac{1}{2}}. \tag{4.7}$$

Now we can determine entropy and pressure as follows:

$$s = \frac{A}{4G} = \frac{(1+\sinh^2\alpha)^{\frac{1}{2}}}{4G}r_0^{(n+1)}\Omega_{n+1}, \tag{4.8}$$

$$P = -\frac{\Omega_{n+1}}{16\pi G}r_0^n, \tag{4.9}$$

$$\varepsilon = Ts - P = (n+1)\frac{\Omega_{n+1}}{16\pi G}r_0^n, \tag{4.10}$$

$$T_{ab} = \frac{\Omega_{n+1}r_0^n}{16\pi G}(nu_a u_b - \eta_{ab}). \tag{4.11}$$

Surprisingly we can see that although the entropy and temperature of charged blackfold differ from neutral blackfold, the concluded stress tensor takes the same form as the neutral blackfold. This proves that expansion of charged and neutral blackfold near horizon takes the same local distribution of energy. It arises from effective description of blackfolds and compactifing of some directions.

5 Conclusion and outlook

Finding the stress energy tensor for all known black holes is very interesting. Depending on the gravity and dimension of spacetime there are different methods that calculate this quantity classically or quantum mechanically. In this paper we calculated the stress energy tensor for higher dimensional black holes which regarded az a black brane which folded into multiple dimensions. This compactified black brane is blackfold. Two cases of blackfols is known: neutral blackfold and charged blackfold. Neutral blackfold has the same metric of extended Schwarzschild black hole but charged one is not Ricci-flat and somewhat a different higher dimensional black hole. We try to find the stress energy tensor for two cases by expansion near horizon. It is the familiar method that uses for spherically symmetric and axisymmetric 4 dimensional black holes. As the calculation shows, the temperature increase by increasing n and depends inversely by the thickness of the horizon. But entropy, pressure, and energy density depend directly to the thickness of horizon.

In the neutral case, as blackfold is Ricci-flat and there is a nice relation between AdS and Ricci-flat which the AdS form of that can be reproduced. It proves that the stress tensor of that AdS form is exactly the stress tensor of the Ricci-flat form by the related transformation. Dependency of temperature to n shows the great influence of topology of blackfold to thermodynamical properties. The different forms of temperature (and also entropy) for two charged and neutral cases, with the same form of stress energy tensor for two cases, proves that local distribution of energy is the same for effective theory of charged and neutral blackfold.

References

[1] Dawood Kothawala, T. Padmanabhan, *Thermodynamic structure of Lanczos-Lovelock field equations from near-horizon symmetries*, Phys.Rev.D79:104020,(2009),[arXiv:0904.0215]

[2] T. Padmanabhan,*Classical and Quantum Thermodynamics of horizons in spherically symmetric spacetimes* Class. Quantum Grav. 19, 5387 (2002) [arXiv:0204019];

[3] T. Jacobson, *Thermodynamics of Spacetime: The Einstein Equation of State* Phys. Rev. Lett. 75, 1260 (1995);

[4] T. Padmanabhan, *Gravity: the inside story* Gen. Rel. Grav. 40, 2031-2036 (2008);

[5] Uma Papnoia, Megan Govenderb, and Sushant G. Ghosha, *Thermodynamic structure of field equations near apparent horizon for radiating black hole*, [arXiv:1411.2323]

[6] J.D. Brown and J.W. York, *Quasilocal energy and conserved charges derived from the gravitational action*, Phys.Rev. D47 1407-1419(1993)[arXiv:9209012]

[7] G. T. Horowitz, *Blackholes in higher dimensions*, Cambridg university press, (2012);

[8] V. Balasubramanian, P. Kraus, *A Stress Tensor For Anti-de Sitter Gravity*, Commun.Math.Phys. 208, 413-428(1999), [arXiv:9902121]

[9] J. M. Maldacena, *The Large N limit of superconformal field theories and supergravity*, Adv. Theor. Math. Phys. 2, 231 (1998), [arXiv:9711200].

[10] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, *Gauge theory correlators from noncritical string theory*, Phys. Lett. B 428, 105 (1998), [arXiv:9802109].

[11] K. Skenderis, *Lecture notes on holographic renormalization*, Class. Quant. Grav. 19, 5849 (2002), [arXiv:0209067].

[12] J. L. Cardy, *Conformal Invariance and Surface Critical Behavior*, Nucl. Phys.B 240, 514 (1984).

[13] A. Bagchi, *Correspondence between Asymptotically Flat Spacetimes and Nonrelativistic Conformal Field Theories*, Phys. Rev. Lett. 105, 171601 (2010).

[14] L. Susskind, âĂIJHolography in the flat space limit,âĂİ AIP Conf.Proc. 493 98(1999) arXiv:hep-th/9901079

[15] A. Bagchi, *The BMS/GCA correspondence*, Phys.Rev.Lett.105:171601(2010), [arXiv:1006.3354].

[16] Marco M. Caldarelli, Joan Camps, Blaise Goutraux, Kostas Skenderis, *AdS/Ricci flat correspondence and the Gregory-Laamme instability* , Phys. Rev. D 87, 061502(R)(2013), [arXiv:1211.2815].

[17] Marco M. Caldarelli, Joan Camps, Blaise Goutraux, Kostas Skenderis, *AdS/Ricci flat correspondence* , JHEP 04 071 (2014), [arXiv:1312.7874].

[18] R. Emparan, T. Harmark, V. Niarchos, N. A. Obers, *World-Volume Effective Theory for Higher-Dimensional Black Holes*, Phys. Rev. Lett. 102 191301 (2009), [arXiv:0902.0427].

[19] R. Emparan, T. Harmark, V. Niarchos, N. A. Obers, *Essentials of Blackfold Dynamics*, JHEP 1003 063(2010), [arXiv:0910.1601].

[20] B. Carter,*Essentials of classical brane dynamics*, Int. J. Theor. Phys. 40 2099-2130(2001), [arXiv:0012036].

[21] R. Emparan, Blackfolds, [arXiv:1106.2021].

[22] J. Sadeghi, Z. Amoozad, *Notes on stress energy tensor of neutral blackfolds*, [arXiv:1504.05370]

[23] Leonard Susskind, James Lindesay, *An Introduction to Black holes, Information and the string Theory Revolution: the Holographic Universe*, World Scientific Publishing (2005)

[24] R. C. Myers, *Stress tensors and Casimir energies in the AdS / CFT correspondence*, Phys. Rev. D 60 046002(1999), [arXiv:9903203]